

**NEWS****OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN  
SERIES CHEMISTRY AND TECHNOLOGY**

ISSN 2224-5286

<https://doi.org/10.32014/2019.2518-1491.69>

Volume 6, Number 438 (2019), 21 – 29

UDC 547.587.11

**Zh.B. Satpaeva<sup>1,2</sup>, O.A. Nurkenov<sup>2</sup>, K.M. Turdybekov<sup>1</sup>, L.K. Abulyaissova<sup>1</sup>,  
M.Zh. Burkeev<sup>1</sup>, S.D. Fazylov<sup>1,2</sup>, S.A. Talipov<sup>3</sup>, David Havlicek<sup>4</sup>**

<sup>1</sup>Ye.A. Buketov Karaganda State University, Kazakhstan;

<sup>2</sup>Institute of Organic Synthesis and Coal Chemistry of the Republic of Kazakhstan, Karaganda;

<sup>3</sup>A.S. Sadykov Institute of Bioorganic Chemistry of the Academy of Sciences  
of Republic Uzbekistan, Tashkent, Uzbekistan;

<sup>4</sup>Charles University, Czech Republic  
(e-mail: [satpaeva\\_zh@mail.ru](mailto:satpaeva_zh@mail.ru), [xray-phyto@yandex.kz](mailto:xray-phyto@yandex.kz))

**MOLECULAR STRUCTURE AND QUANTUM  
CHEMICAL CALCULATIONS  
4-ETHYL-5-(2-HYDROXYPHENYL)-1,2,4-TRIAZOL-3-THIONE**

**Abstract.** The article is devoted to the study of the spatial structure of 4-ethyl-5-(2-hydroxyphenyl)-1,2,4-triazole-3-thione. The molecular and crystalline structure was determined by x-ray diffraction analysis. Data on the spatial structure and crystal packing of the molecule are presented. It is established that phenyl and triazole rings are flat. Due to hydrogen bonds, the molecules in the crystal form three-dimensional networks. The results of an X-ray diffraction study were deposited at the Cambridge Center for Crystal Structural Data. The structure of the synthesized 1,2,4-triazole was studied by <sup>1</sup>H NMR spectroscopy. When analyzing the <sup>1</sup>H NMR spectrum of the compound, characteristic signals of the protons of the aromatic ring are observed. The value of the chemical shift and the integrated signal intensity are determined. A quantum chemical study of 4-ethyl-5-(2-hydroxyphenyl)-1,2,4-triazole-3-thione was carried out by the DFT method using the B3LYP exchange-correlation functional in combination with the Dunning basic set cc-pVDZ. The molecular characteristics of the compound, such as total electron energy, rotational constants, dipole moment and contributions, and thermodynamic functions, are predicted. The equilibrium geometric parameters of the 4-ethyl-5-(2-hydroxyphenyl)-1,2,4-triazole-3-thione molecule were determined. The performed analysis of the spatial configuration and molecular parameters showed a qualitative correspondence between the crystalline and gas-phase structures of the molecule, while it was noted that the main difference is observed in the relative orientation of six- and five-membered cycles.

**Keywords:** crystal structure, x-ray diffraction, 1,2,4-triazole, 4-ethyl-5-(2-hydroxyphenyl)-1,2,4-triazole-3-thione, quantum chemical calculations, DFT method, geometric and energy characteristics.

### **Introduction**

Derivatives of 1,2,4-triazole provide great synthetic possibilities for obtaining new effective drugs. A number of 1,2,4-triazole derivatives have been successfully used in medicine as medicines for the treatment of fungal and viral infections, diseases of the cardiovascular system, etc. [1-4]. New derivatives of 1,2,4-triazole are known, which have shown antiviral [5, 6], anti-inflammatory [7, 8], anticonvulsant [9, 10], antitumor [10, 11], antibacterial [12, 13] and other types of activity. Consequently, some of the 1,2,4-triazole derivatives have been approved as drugs, such as anastrazole, letrozole, risatriptan, ribavirin, alprazolam, fluconazole and posaconazole. This leads to intensive studies of both the preparation methods and the properties of the new derivatives of 1,2,4-triazoles.

It is well known that s-triazoles can exist in two tautomeric forms, and most studies have shown that the thion form was dominant in the solid state and in a neutral solution [14–16]. However, there are some studies that describe the presence of a thiol form [17–20].

Nevertheless, it was shown that thiolthionetautomerism plays an important role in several processes related to biological activity, such as proton transfer and hydrogen bonds [21-24].

Currently, X-ray diffraction analysis is one of the most informative, unique and accurate tools for studying the spatial structure of synthesized compounds, which has made a significant and decisive contribution to the study of almost all inorganic and organic classes [25].

Previously, we synthesized a number of 1,2,4-triazoles based on hydrazides of o- and p-hydroxybenzoic acids, which were obtained by interaction with thiosemicarbazides in an aqueous solution of caustic potassium followed by acidification with acetic acid. X-ray diffraction analysis was used to characterize the crystal structure of 4-allyl-3-(4-hydroxyphenyl)-1H-1,2,4-triazole-5(4H)-thione [26-28]. In continuation of studies in this direction, we present experimental data on the spatial structure and calculated quantum chemical parameters of 4-ethyl-5-(2-hydroxyphenyl)-1,2,4-triazole-3-thione (1).

### Experimental part

The NMR<sup>1</sup>H spectrum was recorded on a Bruker DRX500 spectrometer (500 MHz) in a DMSO-d<sub>6</sub> solution relative to the internal standard — TMS.

**X-ray diffraction analysis** of 4-ethyl-5-(2-hydroxyphenyl)-1,2,4-triazole-3-thione (1). Crystals for X-ray diffraction analysis were obtained by double crystallization of compound 1 from 2-propanol.

Cell parameters and intensities of 3461 reflections (1340 independent, R<sub>int</sub> = 0.0223) were measured on a «Bruker P4» diffractometer (MoK, graphite monochromator, ω scanning, 2.96° ≤ θ ≤ 22.44°) at a temperature of 298 K. Monoclinic crystals, a=9.8479(6), b=7.9892(7), c=13.914(1) Å, β=102.616(3)°, V=1068.3(1) Å<sup>3</sup>, Z=4 C<sub>14</sub>H<sub>9</sub>N<sub>3</sub>O<sub>3</sub>, space group P2<sub>1</sub>/c, d<sub>БНЧ</sub>=1.376 g/cm<sup>3</sup>, μ=0.279 mm<sup>-1</sup>. The initial array of the measured intensities was processed and absorption was taken into account using the SAINT and SADABS programs (multi-scan, T<sub>min</sub> = 0.7072, T<sub>max</sub> = 0.7448) [29].

The structure is deciphered by the direct method. The positions of non-hydrogen atoms are refined in the anisotropic approximation by full-matrix least squares. Hydrogen atoms of the hydroxyl and amino groups were identified from difference synthesis and their positions were refined in the isotropic approximation, the remaining hydrogen atoms were placed in geometrically calculated positions and their positions were refined in the isotropic approximation with fixed positional and thermal parameters (rider model). In the calculations, 1191 reflections of independent reflections with I ≥ 2σ(I) were used, the number of refined parameters is 145. Final divergence factors: R<sub>1</sub> 0.0301, wR<sub>2</sub> 0.0786 (for reflections with I ≥ 2σ(I)), R<sub>1</sub> 0.0347, wR<sub>2</sub> 0.0833 (for all reflections), GooF=1.057. Peaks of residual density: Δρ=0.146 and -0.219 e/Å<sup>3</sup>. The structure has been deciphered and refined using the SHELXS [30] and SHELXL-2018/3 [31] programs. CIF data in the form of a CIF file was deposited at the Cambridge Center for Crystal Structural Data (CCDC 1921694).

**The quantum chemical calculation** for the 4-ethyl-5-(2-hydroxyphenyl)-1,2,4-triazole-3-thione molecule in the gas phase was carried out using the density functional theory method (DFT/B3LYP) [32] using the Gaussian program [33]. In this case, the correlation-consistent Dunning basis set cc-pVDZ was used, which was specially developed to more fully take into account correlation effects.

### Results and discussions

When analyzing the <sup>1</sup>H NMR spectrum of compound 1, characteristic signals of the protons of the aromatic ring are observed. So, the signals of the protons of the aromatic ring are recorded at 7.78 ppm. (J<sub>HαHβ</sub> 8.7 Hz), 6.81 ppm (J<sub>HαHβ</sub> 8.7 Hz), 7.32 ppm (1H, CH<sup>4</sup><sub>arom</sub>), 7.02 ppm. (1H, CH<sup>1</sup><sub>arom</sub>) in the form of doublets, respectively. At 6.95 ppm (1H, CH<sup>2</sup><sub>arom</sub>), 7.42 ppm. (1H, CH<sup>3</sup><sub>arom</sub>) signals appear in the form of triplets. In a weak field, signals of protons of aromatic hydroxyl are recorded at 10.33 ppm. and thioamide N-H proton at 13.8 ppm in the form of two small broadened singlets. Signals of methyl group protons resonate at 1.05 ppm. (J<sub>HH</sub> 7.1 Hz) as a triplet and methylene group at 3.85 ppm. (J<sub>HH</sub> 7.2 Hz) as a quartet.

The spatial structure and packaging of the 4-ethyl-5-(2-hydroxyphenyl)-1,2,4-triazole-3-thione molecule (1) are shown in Figures 1 and 2.




Figure 1 – The molecular structure of the molecule 1




Figure 2 – Packing molecule 1 in the crystal

From the obtained data of X-ray diffraction analysis, it follows that the bond lengths (table 1) and bond angles (table 2) in compounds 1 are close to usual [34]. In molecule 1, the triazole ring is flat; the atoms N1, N2, C3, N4, and C5 are coplanar with an accuracy of  $\pm 0.003 \text{ \AA}$ . The sulfur atom S1, C6 of the ethyl and C8 phenyl groups slightly leave this plane (by -0.051, 0.053 and -0.035  $\text{\AA}$ , respectively). The phenyl cycle, flat with an accuracy of  $\pm 0.001 \text{ \AA}$ , is unfolded relative to the triazole by  $85.6^\circ$ . The O1 atom of the hydroxyl group is practically located in this plane (0.013  $\text{\AA}$ ).

In the crystal, molecules 1 form an intermolecular hydrogen bond N2  $\square$  H ... S1 (1-x, -y, 1 - z) (distance N ... S 3.299 (2)  $\text{\AA}$ , H ... S 2.44 (2)  $\text{\AA}$ , angle N-H ... S 176 (2) $^\circ$ ), linking the triazole moieties of two molecules into dimers. The hydroxyl group forms a hydrogen bond with the N1 atom of another molecule, O1  $\square$  H ... N1 (1-x, 1/2 - y, 1/2 - z) (distance O ... N 2.853(2)  $\text{\AA}$ , H ... N 2.07(2) E, angle O – H ... N 175(2) $^\circ$ ), forming in three-dimensional infinite grids.

A theoretical quantum-chemical approach was used to determine the physicochemical characteristics of 4-ethyl-5-(2-hydroxyphenyl)-1,2,4-triazole-3-thione. In order to find the minimum energy configuration of the molecule in the ground state, the structure is optimized for all independent coordinates. Figure 3 shows an optimized three-dimensional model of the molecule under study (according to quantum chemical calculations). The calculated quantum-chemical structural parameters of the molecule are given (table 1 and table 2) in comparison with the experimental X-ray diffraction data.




Figure 3 – Optimized Molecule 1 Structure (B3LYP/cc-pVDZ Method)

A comparison of the crystalline and gas-phases structures of the molecules showed a qualitative correspondence between them. The main difference is the relative orientation of the two cycles. In the crystalline structure, the torsion angle is  $-93.6^\circ$ , for a molecule in the gas phase it is  $-54.62^\circ$ . According to quantum-chemical calculations, valence bonds between atoms in a gas-phase molecule are more stretched (by 0.001-0.106  $\text{\AA}$ ). Only for N(1)-N(2) and N(4)-C(6) are shortened bonds observed. In the case of O-H and C-H bonds, the calculations give closest to standard (0.96 and 1.09  $\text{\AA}$ , respectively) values (table 1), while P-C bonds are shorter.

Table 1 - Bond lengths in the molecule of 1

Bond	Length, $\text{\AA}$		Angle	Value, deg	
	Exp.	Theor. (B3LYP/cc- pVDZ)		Exp.	Theor. (B3LYP/cc- pVDZ)
S(1)-C(3)	1.6746(19)	1.677	C(3)-N(4)-C(6)	125.18(15)	123.26
O(1)-C(9)	1.358(2)	1.364	C(5)-N(4)-C(6)	126.64(15)	128.39
O(1)-H(01)	0.79(2)	0.969	N(1)-C(5)-N(4)	110.83(15)	111.23
N(1)-C(5)	1.302(2)	1.310	N(1)-C(5)-C(8)	125.76(16)	122.29
N(1)-N(2)	1.378(2)	1.364	N(4)-C(5)-C(8)	123.37(16)	126.32
N(2)-C(3)	1.333(3)	1.362	N(4)-C(6)-C(7)	112.31(17)	113.59

Angle	$\tau$ , deg		Angle	$\tau$ , deg	
	Exp.	Theor. (B3LYP/cc- pVDZ)		Exp.	Theor. (B3LYP/cc- pVDZ)
N(2)-H(2)	0.87(2)	1.012	N(4)-C(6)-H(6A)	109.1	106.78
C(3)-N(4)	1.371(2)	1.388	C(7)-C(6)-H(6A)	109.1	111.52
N(4)-C(5)	1.375(2)	1.390	N(4)-C(6)-H(6B)	109.1	106.32
N(4)-C(6)	1.468(2)	1.466	C(7)-C(6)-H(6B)	109.1	109.92
C(5)-C(8)	1.477(2)	1.478	H(6A)-C(6)-H(6B)	107.9	108.44
C(6)-C(7)	1.502(3)	1.527	C(6)-C(7)-H(7A)	109.5	110.89
C(6)-H(6A)	0.9700	1.099	C(6)-C(7)-H(7B)	109.5	110.22
C(6)-H(6B)	0.9700	1.101	H(7A)-C(7)-H(7B)	109.5	109.46
C(7)-H(7A)	0.9600	1.098	C(6)-C(7)-H(7C)	109.5	109.15
C(7)-H(7B)	0.9600	1.099	H(7A)-C(7)-H(7C)	109.5	108.77
C(7)-H(7C)	0.9600	1.102	H(7B)-C(7)-H(7C)	109.5	108.29
C(8)-C(13)	1.389(3)	1.404	C(13)-C(8)-C(9)	119.56(17)	118.57
C(8)-C(9)	1.390(3)	1.410	C(13)-C(8)-C(5)	120.09(16)	118.88
C(9)-C(10)	1.383(3)	1.400	C(9)-C(8)-C(5)	120.34(16)	122.45
C(10)-C(11)	1.376(3)	1.396	O(1)-C(9)-C(10)	123.47(17)	122.20
C(10)-H(10)	0.9300	1.094	O(1)-C(9)-C(8)	117.03(16)	117.66
C(11)-C(12)	1.367(3)	1.397	C(10)-C(9)-C(8)	119.50(17)	120.14
C(11)-H(11)	0.9300	1.092	C(11)-C(10)-C(9)	120.15(18)	120.28
C(12)-C(13)	1.382(3)	1.395	C(11)-C(10)-H(10)	119.9	120.34
C(12)-H(12)	0.9300	1.091	C(9)-C(10)-H(10)	119.9	119.37
C(13)-H(13)	0.9300	1.091	C(12)-C(11)-C(10)	120.85(18)	120.25
C(9)-O(1)-H(01)	111.1(16)	109.12	C(12)-C(11)-H(11)	119.6	120.35
C(5)-N(1)-N(2)	104.04(15)	103.95	C(10)-C(11)-H(11)	119.6	119.40
C(3)-N(2)-N(1)	113.35(15)	114.55	C(11)-C(12)-C(13)	119.61(19)	119.36
C(3)-N(2)-H(2)	127.6(14)	124.33	C(11)-C(12)-H(12)	120.2	120.51
N(1)-N(2)-H(2)	118.8(14)	121.11	C(13)-C(12)-H(12)	120.2	120.13
N(2)-C(3)-N(4)	103.64(15)	102.33	C(12)-C(13)-C(8)	120.34(18)	121.40
N(2)-C(3)-S(1)	128.90(14)	128.13	C(12)-C(13)-H(13)	119.8	120.46
N(4)-C(3)-S(1)	127.44(15)	129.51	C(8)-C(13)-H(13)	119.8	118.15
C(3)-N(4)-C(5)	108.13(15)	107.93			

Table 2 - Torsion angles in the molecule of 1

C(5)-N(1)-N(2)-C(3)	-1.0(2)	-0.97	N(1)-C(5)-C(8)-C(13)	-93.6(2)	-54.62
N(1)-N(2)-C(3)-N(4)	0.8(2)	1.46	N(4)-C(5)-C(8)-C(13)	83.8(2)	120.31
N(1)-N(2)-C(3)-S(1)	-177.57(14)	-176.67	N(1)-C(5)-C(8)-C(9)	87.9(2)	121.75
N(2)-C(3)-N(4)-C(5)	-0.29(19)	-1.35	N(4)-C(5)-C(8)-C(9)	-94.7(2)	-63.31
S(1)-C(3)-N(4)-C(5)	178.12(14)	176.75	C(13)-C(8)-C(9)-O(1)	-179.38(18)	177.31
N(2)-C(3)-N(4)-C(6)	177.16(16)	-174.46	C(5)-C(8)-C(9)-O(1)	-0.9(3)	0.93
S(1)-C(3)-N(4)-C(6)	-4.4(3)	3.64	C(13)-C(8)-C(9)-C(10)	0.4(3)	-1.68
N(2)-N(1)-C(5)-N(4)	0.78(19)	0.01	C(5)-C(8)-C(9)-C(10)	178.84(18)	-178.07
N(2)-N(1)-C(5)-C(8)	178.45(16)	175.64	O(1)-C(9)-C(10)-C(11)	179.5(2)	-178.02
C(3)-N(4)-C(5)-N(1)	-0.3(2)	0.88	C(8)-C(9)-C(10)-C(11)	-0.2(3)	0.92
C(6)-N(4)-C(5)-N(1)	-177.73(17)	173.53	C(9)-C(10)-C(11)-C(12)	-0.1(3)	0.20
C(3)-N(4)-C(5)-C(8)	-178.07(16)	-174.52	C(10)-C(11)-C(12)-C(13)	0.3(3)	-0.52
C(6)-N(4)-C(5)-C(8)	4.5(3)	-1.88	C(11)-C(12)-C(13)-C(8)	-0.2(3)	-0.27
C(3)-N(4)-C(6)-C(7)	-88.1(2)	-77.10	C(9)-C(8)-C(13)-C(12)	-0.2(3)	1.37
C(5)-N(4)-C(6)-C(7)	88.8(2)	111.27	C(5)-C(8)-C(13)-C(12)	-178.65(19)	177.89

The deviation of the valence angles from similar experimental ranges from 0.04–3.47°, both in the direction of decreasing and increasing values. For dihedral angles, a larger scatter of values is observed when comparing theoretical and experimental data. Differences in the geometric configuration of the 4-ethyl-5-(2-hydroxyphenyl)-1,2,4-triazole-3-thione molecule in different states of aggregation were expected, because, as you know, the geometry of the molecule in the crystal is determined by packing, in pairs - corresponds to the global minimum of total energy.

Theoretical physico-chemical characteristics, such as total energy, zero-point energy, rotational constants, electric dipole moment and others, are given in table. 3.

Table 3 - Physicochemical parameters of molecule 1 according to calculations by the B3LYP/cc-pVDZ method

Options	B3LYP/ cc-pVDZ
Total energy, a.u.	-1025.417093
Rotational Constants, GHz	
A	
B	1.1483336
C	0.3421134 0.2921099
Dipole moment and its components, D	
$\mu_{\text{total}}$	6.265
$\mu_x$	-6.0563
$\mu_y$	1.5974
$\mu_z$	0.1367
$E_{\text{B3MO}}$ , a.e.	-0.19583
$E_{\text{HCMO}}$ , a.e.	-0.04063
$\Delta E_{\text{B3MO-HCMO}}$ , eV	4.223
I, eV	5.329
A, eV	1.106
$\chi$ , eV	3.218
$\eta$ , eV	2.112
S, eV	0.473
$C_v$ , cal/mol·K	51.489
S, cal/mol·K	116.640

Due to the different values of the rotational constants, the molecule is an asymmetric top. Such fundamental properties as the dipole moment, ionization potential, electron affinity, etc., are directly dependent on the geometry of the molecule (table 3). The 4-ethyl-5-(2-hydroxyphenyl)-1,2,4-triazole-3-thione molecule is highly polar. The contributions of the X, Y, and Z components to the total dipole moment are different; the vector of the dipole moment is directed mainly along the X axis.

The most significant orbitals in a molecule are the so-called boundary orbitals: the highest occupied molecular orbital (HOMO) and the lowest free molecular orbital (LUMO). The HOMO energy characterizes the ability to donate an electron, LUMO — to accept an electron, the energy gap between them — the chemical stability of the molecule, with  $I = -E_{\text{VBMO}}$  and  $A = -E_{\text{NCMO}}$  (table 3) [35]. The molecule has a rather large ionization potential, electronegativity, and positive electron affinity. The width of the HOMO-LUMO energy gap ( $\Delta E$ ) indicates a sufficiently low polarizability of the molecule, which is consistent with the large value of its constant dipole moment. The high value of the dipole moment is due to the presence of electronegative atoms in the molecule.

Using these data, the molecular characteristics presented in table 3 were theoretically calculated, such as electronegativity  $\chi$ , chemical rigidity  $\eta$ , and chemical softness S [35]. Absolute electronegativity is calculated as the half-sum of the ionization potential and electron affinity, and absolute chemical rigidity as their half-difference. The stiffness of a substance is considered as a measure of resistance to changes in electronic configuration. Chemical softness is the inverse of stiffness. The table also shows the calculated values of the thermodynamic properties of the compound, such as isochoric heat capacity and entropy.

ӨОЖ 539.19 +548 +547:541.61

**Ж.Б. Сәтбаева<sup>1,2\*</sup>, О.А. Нұркенов<sup>2</sup>, К.М. Тұрдыбеков<sup>1</sup>, Л.К. Абуляисова<sup>1</sup>,  
М.Ж. Бүркеев<sup>1</sup>, С.Д. Фазылов<sup>1,2</sup>, С.А. Талипов<sup>3</sup>, David Havlicek<sup>4</sup>**

<sup>1</sup>Бекетов атындағы Қарағанды мемлекеттік университеті, Қазақстан;

<sup>2</sup>ҚР Органикалық синтез және көмір химиясы институты, Қарағанды, Қазақстан;

<sup>3</sup>ОFA А.С. Садыков атындағы биорганикалық химия институты, Ташкент, Өзбекстан;

<sup>4</sup>Карлов университеті, Чех Республикасы

## **4-ЭТИЛ-5-(2-ГИДРОКСИФЕНИЛ)-1,2,4-ТРИАЗОЛ-3-ТИОННЫҢ МОЛЕКУЛАЛЫҚ ҚҰРЫЛЫМЫ МЕН КВАНТТЫ-ХИМИЯЛЫҚ ЕСЕПТЕУЛЕРИ**

**Аннотация.** Мақала 4-этил-5-(2-гидроксифенил)-1,2,4-триазол-3-тионның кеңістіктік құрылымын зерттеуге арналған. Қосылыстың молекулалық құрылымы рентген құрылымды әдіс арқылы анықталынды. Молекуланың кристалдық қаптамасы мен құрылымы бойынша мәліметтер келтірілген. Фенилді мен триазолды циклдары жазықтықты екені көрсетілген. Молекуланың сұтектік байланысы арқылы кристалда үш өлшемді тордың болатындығы көрсетілген. Рентген құрылымды зерттеу нәтижелері Кембриджтің кристалл құрылымды мәліметтер орталығында тіркелген. Синтезделген 1,2,4-триазолдың құрылымы ЯМР <sup>1</sup>Н-спектроскопия әдісімен зерттелінді. Қосылыстың ЯМР <sup>1</sup>Н спектрінде ароматтық сақинаға тән келетін протондардың сигналы байкалды. Химиялық ығысадың мәні және интегралды сигнал карқындылығы анықталынды. 4-Этил-5-(2-гидроксифенил)-1,2,4-триазол-3-тионды кванттық химиялық өрттей DFT әдісімен B3LYP алмасу-корреляциялық функциясын қолданып, Даннингтүң негізгі жиынтығы cc-pVQZ жиынтығымен жүргізілді. Толық электронды энергия, айналу константалары, диполь моменті мен қосындылар және термодинамикалық функциялар сияқты қосылыстың молекулалық сипаттамалары болжамдалған. 4-Этил-5-(2-гидроксифенил)-1,2,4-триазол-3-тион молекуласының тепе-тес геометриялық параметрлері анықталды. Кеңістіктік конфигурация мен молекулалық параметрлердің талдауы молекуланың кристалды және газды фазалық құрылымдарының арасындағы сапалы сойкестікті көрсетілді, алайда басты айырмашылық алты және бес мүшелі циклдердің салыстырмалы бағытында байқалатындығы анықталынды.

**Түйін сөздер:** кристалдық құрылым, рентген құрылымды анализ, 1,2,4-триазол, 4-этил-5-(2-гидроксифенил)-1,2,4-триазол-3-тион, квантты-химиялық есептеу, DFT әдісі, геометриялық және энергетикалық сипаттамалар.

УДК 539.19 +548 +547:541.61

**Ж.Б. Сатпаева<sup>1,2\*</sup>, О.А. Нұркенов<sup>2</sup>, К.М. Турдыбеков<sup>1</sup>, Л.К. Абуляисова<sup>1</sup>,  
М.Ж. Бүркеев<sup>1</sup>, С.Д. Фазылов<sup>1,2</sup>, С.А. Талипов<sup>3</sup>, David Havlicek<sup>4</sup>**

<sup>1</sup>Карагандинский государственный университет им. Е.А. Букетова, Казахстан;

<sup>2</sup>Институт органического синтеза и углехимии РК, Караганда, Казахстан;

<sup>3</sup>Институт биорганической химии им. Академика Садыкова, Ташкент, Узбекистан;

<sup>4</sup>Карлов университет, Чешская республика

## **МОЛЕКУЛЯРНАЯ СТРУКТУРА И КВАНТОВО-ХИМИЧЕСКИЕ РАСЧЕТЫ 4-ЭТИЛ-5-(2-ГИДРОКСИФЕНИЛ)-1,2,4-ТРИАЗОЛ-3-ТИОНА**

**Аннотация.**Статья посвящена исследованию пространственной структуры 4-этил-5-(2-гидроксифенил)-1,2,4-триазол-3-тиона. Молекулярная и кристаллическая структура определена методом рентгеноструктурного анализа. Приведены данные по пространственному строению и кристаллической упаковке молекулы. Установлено, что фенильные и триазольные циклы плоские. За счет водородных связей молекулы в кристалле образуют трехмерные сетки. Результаты рентгеноструктурного исследования депонированы в Кембриджском центре кристаллоструктурных данных. Исследовано строение синтезированного 1,2,4-триазола методом ЯМР <sup>1</sup>Н-спектроскопии. При анализе спектра ЯМР <sup>1</sup>Н соединения наблюдаются характерные сигналы протонов ароматического кольца. Определено значение химического сдвига и интегральная интенсивность сигналов. Методом DFT с использованием обменно-корреляционного функционала B3LYP в сочетании с базисным набором Даннингасс-рVQZ проведено квантово-химическое исследование 4-этил-5-(2-гидроксифенил)-1,2,4-триазол-3-тиона. Предсказаны молекулярные характеристики соединения, такие как полная электронная энергия, вращательные постоянные, дипольный момент и

вклады, термодинамические функции. Определены равновесные геометрические параметры молекулы 4-этил-5-(2-гидроксифенил)-1,2,4-триазол-3-тиона. Выполненный анализ пространственной конфигурации и молекулярных параметров показал качественное соответствие кристаллической и газофазной структур молекулы, при этом отмечено, что основное отличие наблюдается в относительной ориентации шести- и пятичлененных циклов.

**Ключевые слова:** кристаллическая структура, рентгеноструктурный анализ, 1,2,4-триазол, 4-этил-5-(2-гидроксифенил)-1,2,4-триазол-3-тион, квантово-химические расчеты, метод DFT, геометрические и энергетические характеристики.

**Information about the authors:**

Satpaeva Zh.B. – Doctoral student of Buketov Karaganda State University, e-mail: [satpaeva\\_zh@mail.ru](mailto:satpaeva_zh@mail.ru), <https://publons.com/researcher/P-4887-2017/>, <https://orcid.org/0000-0003-0962-1148>

Nurkenov O.A. – Doctor of chemical sciences, Professor, Institute of Organic Synthesis and Coal Chemistry of the Republic of Kazakhstan, e-mail: [nurkenov\\_oral@mail.ru](mailto:nurkenov_oral@mail.ru), <https://orcid.org/0000-0002-2771-0411>

Turdybekov K.M. – Doctor of chemical sciences, Professor, Department of Physics and Nanotechnology of Buketov Karaganda State University, e-mail: <https://orcid.org/0000-0001-9625-0060>

Abulyassova L.K. – Candidate of Chemical Sciences, Professor of the Department of Physical and Analytical Chemistry of Buketov Karaganda State University, e-mail: [lyazzat.kz07@mail.ru](mailto:lyazzat.kz07@mail.ru), <https://orcid.org/0000-0002-7530-3378>

Burkeev M.Zh. – Acting Dean of the Faculty of Chemistry of Buketov Karaganda State University, e-mail: [m\\_burkeev@mail.ru](mailto:m_burkeev@mail.ru), <https://orcid.org/0000-0001-8084-4825>, <https://publons.com/researcher/2339931/meyram-z-burkeev/>

Fazylov S.D. – corresponding member of NAS RK, Doctor of chemical sciences, Professor, Institute of Organic Synthesis and Coal Chemistry of the Republic of Kazakhstan. e-mail: [iosu8990@mail.ru](mailto:iosu8990@mail.ru), <https://orcid.org/0000-0002-4240-6450>

Talipov S.A. – A.S. Sadikov Institute of Bioorganic Chemistry, Uzbekistan Academy of Science, e-mail: [samat\\_talipov@yahoo.com](mailto:samat_talipov@yahoo.com)

David Havlicek – Department of Inorganic Chemistry, Charles University in Prague. <https://orcid.org/0000-0002-8854-6213>; <https://publons.com/researcher/B-1376-2017/>; Scopus Author ID 6602118880

## REFERENCES

- [1] Hassan H.Y., El-Shorbagi A.A., El-Koussi N.A., Abdel-Zaher A.O. (1994) Desing and synthesis of some new 1H-1,2,4-triazoles of potential anti-inflammatory and analgesic activities, Bulletin of Pharmaceutical Sciences, 17:27-39. (in Eng)
- [2] Turan-Zitouni G., Kaplancikli Z.A., Erol K., Kilic F.S. (1999) Synthesis and analgesis activity of some triazoles and triazolothiadiazines, Farmaco, 54:218-223.
- [3] Kokil G.R., Rewatkar P.V., Gosain S., Aggarwal S., Verma A., Karla A., et al. (2010) Synthesis and in vitro evaluation of novel 1,2,4-triazole derivatives as antifungal agents, Letters of Drug Desing and Discovery, 7:46-49. DOI: 10.2174/157018010789869415 (in Eng).
- [4] Turan-Zitouni G., Kaplancikli Z.A., Yildiz M.T., Chevallet P., Kaya D. (2005) Synthesis and antimicrobial activity of 4-phenyl/cyclohexyl-5-(1-phenoxyethyl)-3-[N-(2-thiazolyl)acetamido]-thio-4H-1,2,4-triazole derivatives, European Journal of Medicinal Chemistry, 40:607-613. DOI: 10.1016/j.ejmech.2005.01.007 (in Eng).
- [5] Kritsanida M., Mouroutsou A., Marakos P., Pouli N., Papakonstantinou-Garaufalias S.M., Pannecouque C., et al. (2002) Synthesis and antiviral activity evalution of some new 6-substituted 3-(1-adamantyl)-1,2,4-triazolo[3,4-b][1,3,4]thiadiazoles, Farmaco, 57:253-257. PMID: 11989804 (in Eng).
- [6] Johns B.A., Weatherhead J.G., Allen S.H., Thompson J.B., Garvey E.P., Foster S.A., et al. (2009) The use of axadiazole and triazole substituted naphthyridines as HIV-1 integrase inhibitors. Part 1: Establishing the pharmacophore, Bioorganical and Medicinal Chemistry Letters, 19:1802-1806. DOI: 10.1016/j.bmcl.2009.01.090. PMID: 19217781 (in Eng).
- [7] Mathew V., Keshavayya J., Vaidya V.P., Giles D. (2007) Studies on synthesis and pharmacological activities of 3,6-disubstituted-1,2,4-triazolo[3,4-b]-1,3,4-thiadiazoles and their dihydro analogues, European Journal of Medicinal Chemistry, 42(6):823-840. DOI: 10.1016/j.ejmech.2006.12.010 (in Eng).
- [8] Tozkoparan B., Kupeli E., Yesilada E., Ertan M. (2007) Preparation of 5-aryl-3-alkylthio-1,2,4-triazoles and corresponding sulfones with anti-inflammatory-analgesic activity, Biorganical and Medicinal Chemistry, 15:1808-1814. DOI: 10.1016/j.bmc.2006.11.029. PMID: 17166724 (in Eng).
- [9] Kelley J.L., Koble C.S., Davis R.G., McLean E.W., Soroko F.E., Cooper B.R. (1995) 1-(Fluorobenzyl)-4-amino-1H-1,2,3-triazolo[4,5-c]pyridines: Synthesis and anticonvulsant activity, Journal of Medicinal Chemistry, 38:4131-4134. (in Eng)
- [10] Holla B.S., Poojary K.N., Rao B.S., Shivananda M.K. (2002) New bis-aminomercaptotriazoles and bis-triazolothiadiazoles as possible anticancer agents, European Journal of Medicinal Chemistry, 37(6):511-517. PMID: 12204477 (in Eng).
- [11] Sztanke K., Tuzimski T., Rzymowska J., Pasternak K., Kandefer- Szerszen M. (2008) Synthesis, determination of the lipophilicity, anticancer and antimicrobial properties of some fused 1,2,4-triazole derivatives, European Journal of Medicinal Chemistry, 43:404-419. DOI: 10.1016/j.ejmech.2007.03.033 (in Eng).
- [12] Bay H.A., Quaddouri B., Guadaoui A., Touzani R., Benchat N.E., Hamal A., et al. (2010) Synthesis and biological activity of new triazole compounds, Drug Desing and Discovery, 7:737-742. DOI: 10.2174/157018010789869352 (in Eng).

- [13] Plech T., Wujec M., Kosikowska U., Malm A., Rajtar B., Polz-Dacewicz B. (2013) Synthesis and in vitro activity of 1,2,4-triazole-ciprofloxacin hybrids against drug-susceptible and drug-resistant bacteria, European Journal of Medicinal Chemistry, 6:128-134. DOI: 10.1016/j.ejmech.2012.11.040 (in Eng).
- [14] Roy G., Mugesh G. (2005) Anti-thyroid drugs and thyroid hormone synthesis: Effect of methimazole derivatives on peroxidase-catalyzed reactions, Journal of the American Society, 127:15207-15217. DOI: 10.1021/ja054497u (in Eng).
- [15] Wujec M., Pisklak M., Pitucha M., Dobosz M., Siwek A. (2007) The reactions of hydrazide (4-nitroimidazol-1-yl)acetic acid with isothiocyanates, Annals of the Polish Chemical Society, 604-606. (in Eng).
- [16] Wujec M., Paneth P. (2008) Mechanism of 4-methyl-1,2,4-triazol-3-thione reaction with formaldehyde, Journal of Physical Organic Chemistry, 20:345-348. DOI: 10.1002/poc.1324 (in Eng).
- [17] Plech T., Luszczki J., Wujec M., Flieger J., Pizon M. (2013) Synthesis, characterization and preliminary anticonvulsant evaluation of some 4-alkyl-1,2,4-triazoles, European Journal of Medicinal Chemistry, 60:208-215. DOI: 10.1016/j.ejmech.2012.11.026. PMID: 23291122 (in Eng).
- [18] Roy G., Nethaji M., Mugesh G. (2004) Biomimetic studies on anti-thyroid dregs and thyroid hormone synthesis, Journal of the American Society, 126,9:2712-2713. DOI: 10.1021/ja039860g (in Eng).
- [19] Zamani K., Faghihi K., Tofighi T., Shariatzadeh M.R. (2004) Synthesis and antimicrobial activity of some pyridyl and naphthyl substituted 1,2,4-triazole and 1,3,4-thiadiazole derivatives, Turkish Journal of Chemistry, 28:95-100. (in Eng).
- [20] Mroczeck T., Plech T., Wujec M. (2017) Novel Concept of Discrimination of 1,2,4-triazole-3-thione and 3-thiol tautomers, Journal of Chromatographic Science, 55, 2:117-129. DOI: 10.1093/chromsci/bmw151 (in Eng).
- [21] Coyanis E.M., Della Vedova C.O., Haas A., Winter M. (2002) Preparation, characterization and thiol-thionetautomeric studies of 2-thiono-4-methyl-5-(2,2,2-trifluoro-1-trifluoromethyl ethyl)-1,3-thiazoline, Journal of Fluorescence Chemistry, 117(2):185-192. DOI: 10.1016/S0022-1139(02)00186-0 (in Eng).
- [22] Anuva S., Bijan K.P., Samiran K., Nihil G. (2011) Excited state lactim to lactam type tautomerization reaction in 5-(4-fluorophenyl)-2-hydroxypyridine: Spectroscopic study and quantum chemical calculation, Journal of Fluorescence, 21:95-104. DOI: 10.1007/s10895-010-0692-3. PMID: 20614169 (in Eng).
- [23] Bach A., Tanner C., Manca C., Frey H.M., Leutwyler S. (2003) Ground- and excited state proton transfer and tautomerization in 7-hydroxyquinoline ( $\text{NH}_3$ )<sub>n</sub> clusters: Spectroscopic and time resolved investigations, Journal of Chemical Physics, 119(12):5933-5942. DOI: 10.1063/1.1603740 (in Eng).
- [24] Douhal A., Kim S.K., Zewail A.H. (1995) Femtosecond molecular dynamics of tautomerization in model base pairs, Nature, 378:260-263. (in Eng).
- [25] Bokay G.B., Poray-Koshits M.A. (1964) X-ray analysis. Moskay.
- [26] Nurkenov O.A., Fazylov S.D., Satpaeva Zh.B., Seilkhanov T.M., Karipova G.Zh., Isaeva A.Zh. (2014) Synthesis and structure of new derivatives of salicylic acid hydrazide, Russ. J. Gen. Chem. 84, 9:1857-1859. DOI: 10.1134/S1070363214090369 (in Eng.).
- [27] Nurkenov O.A., Fazylov S.D., Satpaeva Zh.B., Turdybekov K.M., Karipova G.Zh., Isaeva A.Zh., Talipov S.A., Ibragimov B.T. (2015) Synthesis and structure of new 1,2,4 – triazoles derived from *p*-hydroxybenzoic acid hydrazide, Russ. J. Gen. Chem. 85, 1:57-60. DOI: 10.1134/S1070363215010107 (in Eng).
- [28] Nurkenov O.A., Fazylov S.D., Zhivotova T.S., Satpayeva Zh.B., Akhmetova S.B., Kurmanbayeva Zh.Zh., Karipova G.Zh., Isaeva A.Zh. (2015) Synthesis, chemical transformations and antimicrobialactivity of somethiosemicarbazidesof o- and p-hydroxybenzoicacid, News of the National academy of sciences of the Republicof Kazakhstan. Ser. chem. andtech. 1, 409:9-14. (in Eng)
- [29] SMART V5.051 and SAINT V5.00, (1998) Area detector control and integration software, Bruker AXS Inc., Madison, WI-53719, USA.
- [30] Sheldrick G.M. (2008). ActaCrystallographica Section A: Foundations of Crystallography. International Union of Crystallography 64:112.
- [31] Sheldrick G.M. (2015). ActaCrystallographica Section C: 71:3.
- [32] Becke A.D. (1993) J. Chem. Phys., 98, 7:5648.
- [33] Frisch M.J., Trucks G.W., Schlegel H.B. et al. (2016) Gaussian 16, Revision A.03, Wallingford CT.
- [34] Allen F.H., Kennard O., Watson D.G., Brammer L., Orpen A.G., Taylor R. (1987). Tables of bond lengths determined by x-ray and neutron diffraction, JournalChemistry Society Perkin Trans. 2, 1.
- [35] Fleming I. (1976) Frontier Orbitals and Organic Chemical Reactions. Wiley, London. 249.
- [36] KenzhaliyevB. K. et al., Development of technology for chromite concentrate from the slurry tailings of enrichment.News of the National Academy of Sciences of the Republic of Kazakhstan series of geology and technical sciences. Volume 3, Number 429 (2018), 182 – 188.
- [37] Presnyakov, A. A., Kenzhaliiev, B. K., Kozhakhmetov, S. M., Panichkin, A. V., & Ponomareva, S. V. (2002). Metal Science and Heat Treatment, 44(1/2), 80–80. <https://doi.org/10.1023/a:1015341013948>
- [38] Kenzhaliiev, B.K., Tashuta, G.N., Valutskaya, T.A. et al. Potentiometric Determination of Mercury with Iodide-Selective Electrodes. Journal of Analytical Chemistry (2002) 57: 261. <https://doi.org/10.1023/A:1014456602075>