

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

SERIES OF AGRICULTURAL SCIENCES

ISSN 2224-526X

Volume 1, Number 43 (2018), 42 – 44

UDC 616.1/9:636.2(574. 51)

A. A. Jangabulova¹, A. Z. Maulanov², A. A. Zhumageldiev¹

¹Biological Safety Department, Kazakh national agrarian university, Almaty, Kazakhstan,

²Veterinary Sanitary Expertise and Hygiene Department, Kazakh national agrarian university, Almaty, Kazakhstan.

E-mail: ermaz@inbox.ru, ainur_89_jan@mail.ru

**HISTOLOGY OF MEAT PRODUCTS
FROM CATTLE WITH KETOSIS**

Abstract. At present, all livestock dairy farms throughout the world suffer from ketosis. Ketosis mostly affects highly-productive animals. Ketosis causes severe clinical manifestations and sometimes mortality. The productivity of dairy cows with ketosis decreases to the minimum and recovered animals lose their previous productivity.

Key words: ketosis, histology, dystrophia, haemotoxylin eosin, hepatocyte, pathoanatomy.

Introduction. Prevalence of ketosis of dairy cows in the countries of the European Union and in the most widespread control systems of after calving individuals during a transition period till recently remained substantially unexplored, except for the works reporting about the high frequency of occurrence of this pathology and also for the big economic losses of dairy production, related until recently (McKay, 2012).

The recent research conducted from May 2011 till October 2012 in 10 European countries showed that in dairy production the prevalence of cattle's ketosis, defined as a high concentration of β -hydroxyoleic acid in serum of blood in cows was ranging from 21.8 to 36.6% [1-4].

Frequency of occurrence of level of ketosis at individuals in dairy herd was the lowest in the United Kingdom and Italy (31%) and was much higher in Germany (43%), in the Netherlands (46%) and in France (53%). In five of the studied countries the average value of prevalence of ketosis in herd of young dairy cows in the range of observation from 1 to 3 weeks after calving made 41%, while the observation within 35 days after calving gave the frequency of diagnostics of a clinical stage of ketosis only in 1.6% [5-8].

According to the definition [9, 10] ketosis or the acetonemia is the metabolic disorder which is characterized by abnormal increase in ketone bodies in blood of dairy cows which, usually, happens in the postnatal period, between the 2nd and 6th week after calving. Economic factors of production to the great extent suffer from influence of ketosis, because of the decrease in production of milk, because of the costs of treatment of individuals, because of the increased risk of a susceptibility to other diseases, the decrease in efficiency of reproduction and more other risks [11-13].

In clinical aspect of veterinary science a ketosis in dairy cows, is more often described in scientific publications as subclinical ketosis. The main aspects of a research of ketosis already occur in the scientific periodical press from the beginning of 1950ies (Holmes, 1950). Later on, because of the increase in productivity and the expansion of production of dairy products this metabolic disorder became more widespread. This led to the appearance of new scientific works [14-20].

Work objective: histology of meat products from cattle with ketosis.

Methods. Meat products of forcibly killed cows were used for histological examination. The sample was taken from muscle, liver, kidney and heart of the cow. Obtained samples were put into 10% aqueous solution of 10×10×4 mm of formalin and then painted with hematoxylin eosin using standard method. The microscopy was performed using microscope. The stereometric analysis was carried out using the measuring grid.

Results. According to histology, dystrophic changes of different levels were noted in skeletal muscles, liver, kidneys and heart. Protein and fatty degeneration in liver cells have been identified as stable and consistent. Hepatocytes in histological preparations painted with hematoxylin eosin were in high level and their cytoplasm had small, large empty interspaces.

Hepatic volumes were noted as expanded and rounded, and interspaces in cytoplasm composed large volumetric pores. Such interspaces in some hepatocytes occupied completely all cytoplasm and the cell nucleus shifts to the edge of the cytoplasm. Small vessels are depressed and abnormal. As well as, proliferation of reticuloendothelial and mononuclear macrophage system cells has been constantly noted in all histologic preparations. They were found in reticular and lymphoid forms. Fat pieces in painted preparations were painted in red and brown. In the Schiff (PAS) reaction the amount of glycogen has decreased to the minimum in comparison to control group. The skeletal muscles are thickened at different levels, the horizontal strips are not clearly visible, and sarcoplasm had rounded grains and fat pieces painted with eosin in red. Also, the accumulation of fatty tissue in interstitial tissue caused atrophic compression of meat fibers. In the Schiff (PAS) reaction the amount of glycogen has decreased to the minimum in comparison to control group. The horizontal lines of the heart are almost invisible, and sarcoplasm had rounded grains and fat pieces painted with eosin in red. Also, the accumulation of fatty tissue in interstitial tissue caused atrophic compression of meat fibers. In the Schiff (PAS) reaction the amount of glycogen has decreased to the minimum in comparison to control group.

Dicussion. It was found that all the kidneys under test were damaged. These changes were caused by dystrophy and necrobiosis, as well as reflected as parenchymatous and adipose degenerations in epithelial cells of kidney's vertical holes. Some epithelial cells were destroyed by necrobiosis. Infiltration of polymorphic cells was noted in intermediate connective tissue. In the Schiff (PAS) reaction the amount of glycogen has decreased to the minimum in comparison to control group.

Conclusion. According to histology results, internal organs of dairy cows differed in the presence of parenchymatous, adipose and carbohydrate degenerations.

REFERENCES

- [1] Agresti A. 2002a. Analyzing repeated categorical response data. Pages 455–490 in *Categorical Data Analysis*. John Wiley & Sons Inc. New York, NY.
- [2] Agresti A. 2002b. Multinomial response models. Pages 267–302 in *Categorical Data Analysis*. John Wiley & Sons Inc., New York, NY.
- [3] Andersson L. 1988. Subclinical ketosis in dairy cows. *Vet. Clin. North Am. Food Anim. Pract.* 4:233–251.
- [4] Andersson L., Emanuelson U. 1985. An epidemiological study of hyperketonaemia in Swedish dairy cows; determinants and the relation to fertility. *Prev. Vet. Med.* 3:449–462.
- [5] Asl A.N., Nazifi S., Ghasrodashti A.R., Olyae A. 2011. Prevalence of subclinical ketosis in dairy cattle in the South-western Iran and detection of cutoff point for NEFA and glucose concentrations for diagnosis of subclinical ketosis. *Prev. Vet. Med.* 100:38–43.
- [6] Amory J.R., Barker Z.E., Wright J.L., Mason S.A., Blowey R.W., Green L.E. 2008. Associations between sole ulcer, white line disease and digital dermatitis and the milk yield of 1824 dairy cows on 30 dairy cow farms in England and Wales from February 2003–November 2004. *Prev. Vet. Med.* 83:381–391. <http://dx.doi.org/10.1016/j.prevetmed.2007.09.007>.
- [7] Archer S.C., Green M.J., Huxley J.N. 2010. Association between milk yield and serial locomotion score assessments in UK dairy cows. *J. Dairy Sci.* 93:4045–4053. <http://dx.doi.org/10.3168/jds.2010-3062>.
- [8] Alban L. 1995. Lameness in Danish dairy cows: Frequency and possible risk factors. *Prev. Vet. Med.* 22:213–225.
- [9] Bicalho R.C., Warnick L.D., Guard C.L. 2008. Strategies to analyze milk losses caused by diseases with potential incidence throughout the lactation: A lameness example. *J. Dairy Sci.* 91:2653–2661. <http://dx.doi.org/10.3168/jds.2007-0744>.
- [10] Batanova O.V., Jelenshleger A.A. *Morfologicheskie i biohimicheskie pokazateli krovi u korov pri ketoze // Molodezh' Barnaulu*. Barnaul, 2005. P. 316-317.
- [11] Batanova O.V., Jelenshleger A.A. *Lechenie korov, bol'nyh ketozom // Vestnik Altajskogo gosudarstvennogo agrarnogo universiteta*. Barnaul, 2006. P. 40-42.
- [12] Batanova O.V., Jelenshleger A.A. *Profilaktika subklinicheskogo ketoza korov // Vestnik Altajskogo gosudarstvennogo agrarnogo universiteta*. Barnaul, 2006. P. 32-34.
- [13] Batanova O.V., Dutova O.G., Jelenshleger A.A. *Funkcional'noe sostojanie shhitovidnoj zhelezy pri lechenii ketoza korov // Sibirskij Vestnik sel'skohozjajstvennoj nauki*. Krasnoobsk, 2007. P. 66-68.
- [14] Batanova O.V., Jelenshleger A.A. *Tireoidnyj status korov pri ketoze // Aktual'nye problemy sel'skogo hozjajstva gornyh territorij. Gorno-Altajsk*, 2007. N 1. P. 57-59.
- [15] Batanova O.V. *Soderzhanie ketonovyh tel i tireoidnyh gormonov v krovi korov pri ketoze // Veterinarija*. 2008. N 2. P. 43-45.

[16] Baird G.D., 1982. Primary ketosis in the high-producing dairy cow: Clinical and subclinical disorders, treatment, prevention, and outlook. *J. Dairy Sci.* 65:1–10.

[17] Barker Z.E., Amory J.R., Wright J.L., Blowey R.W., Green L.E. 2007. Management factors associated with impaired locomotion in dairy cows in England and Wales. *Dairy J. Sci.* 90:3270–3277.

[18] Barker Z.E., Leach K.A., Whay H.D., Bell N.J., Main C.J. 2010. Assessment of lameness prevalence and associated risk factors in dairy herds in England and Wales. *Dairy J. Sci.* 93:932–941.

[19] Bicalho R.C., Warnick L.D., Guard C.L. 2008. Strategies to analyze milk losses caused by diseases with potential incidence throughout the lactation: A lameness example. *Dairy J. Sci.* 91:2653–2661.

[20] Borchardt S., Staufenbiel R. 2012. Evaluation of the use of nonesterified fatty acids and β -hydroxybutyrate concentrations in pooled serum samples for herd-based detection of subclinical ketosis in dairy cows during the first week after parturition. *Am J. Vet. Med. Assoc.* 240:1003–1011.

А. А. Джангабулова¹, А. З. Мауланов², А. А. Жумагелдиев¹

¹Кафедра «Ветеринариялық-санитариялық сараптау және гигиена»
Қазақ ұлттық аграрлық университеті, Алматы, Қазақстан,

²Кафедра «Биологиялық қауіпсіздік» Қазақ ұлттық аграрлық университеті, Алматы, Қазақстан,

КЕТОЗБЕН АУЫРҒАН ІРІ ҚАРА МАЛ ЕТ ӨНІМДЕРІНІҢ ГИСТОЛОГИЯСЫ

Аннотация. Кәзіргі кезде кетоз әлемнің барлық елдеріндегі сүт бағытындағы малшаруашылықтарында тіркеледі. Кетозға бірінші кезекте жоғары өнім беретін жануарлар шалдығытыны анықталған. Кетоз өте зілді клиникалық белгілермен өтеді және кейде өліммен аяқталатыны байқалған. Ауырған сиырлардың сүт өнімі төмендейді, емделгеннен кейін оның бұрынғы деңгейдегі сүт өнімі қалпыны келмейді.

Түйін сөздер: кетоз, гистология, дистрофия, гематоксилин эозин, гепатоцит, патологоанатомия.

А. А. Джангабулова¹, А. З. Мауланов², А. А. Жумагелдиев¹

¹Кафедра «Ветеринарная-санитарная оценка и гигиена»,
Казахский национальный аграрный университет, Алматы, Казахстан,

²Кафедра «Биологическая безопасность»,
Казахский национальный аграрный университет, Алматы, Казахстан

ГИСТОЛОГИЯ МЯСА ПРИ КЕТОЗЕ

Аннотация. В настоящее время кетоз зарегистрирован в животноводческих молочных фермах во всех странах мира. Кетоз проходит через крайние клинические симптомы и иногда заканчивается смертью. При болезни кетоза уменьшается подукт молока, и после лечения у нее не будет предыдущего молочного продукта.

Ключевые слова: кетоз, гистология, дистрофия, гематоксилин эозин, гепатоцит, патологоанатомия.